



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2019; 7(3): 1347-1350

© 2019 JEZS

Received: 01-03-2019

Accepted: 03-04-2019

Rajat Deshwal

Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

SK Sachan

Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

Gaje Singh

Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

DV Singh

Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

Gopal Singh

Plant Pathology, Sardar
Vallabhbhai Patel University of
Agriculture & Technology
Meerut, Uttar Pradesh, India

Pooran Chand

Genetics & Plant Breeding,
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

Correspondence**Rajat Deshwal**

Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture &
Technology Meerut,
Uttar Pradesh, India

Seasonal abundance of insect pests associated with paddy crop in western plain zone of Uttar Pradesh

Rajat Deshwal, SK Sachan, Gaje Singh, DV Singh, Gopal Singh and Pooran Chand

Abstract

Field experiment was conducted to study the insect pests and their natural enemies associated with basmati rice in western plain zone of Uttar Pradesh during *Kharif* 2017 and 2018 at Crop Research Center of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.), India. During the period of study, twelve insect species were found associated with basmati rice in this region, which belong to six orders *viz.* lepidoptera (yellow stem borer, leaf folder and swarming caterpillar), homoptera (green leaf hopper, brown plant hopper and white backed plant hopper), heteroptera (rice gundhi bug), coleoptera (rice root weevil and white grub), isoptera (termite) and orthoptera (grass hopper). Among natural enemies rove beetles, tiger beetles, carabid beetles, damsel flies and dragon flies were recorded from the rice agro-ecosystem.

Keywords: Insect pests, basmati rice, yellow stem borer, leaf folder and natural enemies

Introduction

Rice is one of the most important staple food crop of India for more than 2/3rd of its population. The slogan "Rice is life" can be considered appropriate for our country as this crop plays a vital role in our national food security and is a means of livelihood for millions of rural households. Rice is one of the world's largest cereal crop providing the caloric need for millions of people. In the World, rice is grown in 167.24 million ha with an annual production of 769.65 million tonnes and 4601.9 kg /ha of productivity and in India area under rice cultivation is 43.78 million ha, production is 168.50 million tones and yield is 3848 kg/ha^[1]. Haryana, Punjab, Uttaranchal, Uttar Pradesh and Jammu Kashmir are the States, where basmati rice is grown in our country. Basmati rice crop suffers severely due to attack of various insect pests, which reduces its yield and quality. More than 100 species of insects have been reported to attacked rice crop from the germination of nursery till its harvests. In general, yield loss due to insect pest of rice has been estimated at about 25% in different rice ecosystem^[10]. Therefore, the detailed studies on insect pests associated with basmati rice have been undertaken.

2. Materials and Methods

The present study was conducted at the Crop Research Centre of Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, U.P., India during *Kharif* 2017 and 2018. Insect-pests of rice and their natural enemies were recorded after transplanting to harvest of the crop at weekly intervals. Observations were taken randomly at ten spots of each plot. The insects were collected and identified. The nature and extent of damage caused by various insect pests was also recorded to assess the economic status of pest.

3. Results and Discussion

During study, twelve insect species belonging to six orders and eight families were recorded on basmati rice at different crop growth stages. Among them, yellow stem borer, *Scirpophaga incertulas* Walker and leaf folder, *Cnaphalocrocis medinalis* Guenee were found as major pests. The brown plant hopper, *Nilaparvata lugens* Stal., white grub, *Holotrichia consanguinea* Blanch, termite, *Odontotermes obesus* Romb. and kharif grass hopper, *Hieroglyphus banian* Fab. were found moderately damaging the crop. The rice swarming caterpillar (army worm),

Spodoptera mauritia Boisduval, green leaf hopper, *Nephotettix virescens* Distant, white backed plant hopper, *Sogatella furcifera* Horvath, rice gundhi bug, *Leptocorisa acuta* Thunb, rice root weevil, *Echinocnemus oryzae* Marshall, and grass hopper, *Oxya fuscovittata* Marshall recorded on the crop were of less importance and extent of their damage was found without much economic loss (Table-1).

Yellow stem borer, *Scirpophaga incertulas* Walker (Pyralidae : Lepidoptera) was recorded as most destructive and major insect pest of rice crop in this region. The appearance of this pest starts in the beginning of July and remained active throughout the crop season. The larvae of this insect bore inside the stem from the growing point and fed internally causing death of central shoot “dead heart” in vegetative stage and “white ear head” at reproductive stage, respectively. This resulted in chaffy grains. The damaged plants were easily pulled out. The severe damage of this pest was recorded from beginning of August to September end. The damage of yellow stem borer on rice crop has also been reported Kumar and Patil (2004), Gowda and Gubbaiah (2011), Kashyap (2013) and Gangwar *et al.* (2015). Sachan *et al.* (2006) also reported the severe incidence of *S. incertulas* on basmati rice throughout the crop season in tarai region of Uttar Pradesh [2-5, 9].

Leaf folder, *Cnaphalocrocis medinalis* Guenee (Pyralidae: Lepidoptera), previously known as a minor foliage feeding pest has acquired the status of major pest of basmati rice and recorded during August–September. The young larvae fed on tender leaves without folding them. The second instars larvae glues the growing paddy leaves longitudinally for accommodation and fed voraciously green foliage, which results in papery dry leaves. Feeding greatly reduced the general vigor and photosynthetic ability of an infested plant. The swarming caterpillar, *Spodoptera mauritia* Boisduval (Noctuidae: lepidoptera) larvae fed on the upper portion of rice canopy by defoliating leaves during night. This insect recorded as minor pest in the month of July–August. Considerable losses to paddy crop due to leaf folder have been reported by several workers like Sachan *et al.* (2006), Gowda and Gubbaiah (2011), Kashyap (2013), Singh and Singh (2014) and Gangwar *et al.* (2015) [2-4, 9, 11].

The leaf hoppers and plant hoppers are sucking insects, which remove plant sap from xylem and phloem tissues of the plant. The incidence of green leaf hopper, *Nephotettix virescens* Distant (Cecadellidae: Homoptera) was recorded during August–September. Both nymphs and adults of this insect sucked the plant sap from the leaves and tender part of plant by turning them yellow. They may also transmit viral disease to the plant. This insect was recorded as minor pest from this region. Brown plant hopper, *Nilaparvata lugens* Stal. (Delphacidae : Homoptera) is another important pest of rice. Its infestation was recorded from middle of August to September end. As a result of feeding by both nymphs and adults at the base of the tillers, plants turn yellow and dry up rapidly. At early infestation round yellow patches appeared which soon turn brownish due to the drying up of the plants. This condition is called “hopper burn”. Complete distraction of the crop was recorded in severe cases. White backed plant hopper, *Sogatella furcifera* Horvath (Delphacidae: Homoptera) has been recorded in August - September as minor pest in this region. It sucked the sap from tender leaves, thus causing yellowishness of them. The honey dew produced by the hoppers serves as a medium for mould growth. The

damage caused by green hopper, brown plant hopper, and white backed plant hopper on rice crop has been reported by various workers such as Sachan *et al.* (2006), Kashyap (2013) and Singh and Singh (2014) [4, 9, 11].

The rice gundhi bug, *Leptocorisa acuta* Thunb. (Coreidae: Heteroptera) was recorded as important pest of rice crop in this region. Both nymphs and adults sucked the sap of grains during milky stage and thus make them chaffy. Whole panicle becomes white colored (chaffy) under severe infestation. Its occurrence was recorded during September–October. Sachan *et al.* (2006), Kashyap (2013), Singh and Singh (2014) and Gangwar *et al.* (2015) also reported the damage of this pest during September–October on rice crop [3, 4, 9, 11].

Rice Root Weevil, *Echinocnemus oryzae* Marshall (Curculionidae: Coleoptera) was also recorded a common pest of basmati rice at vegetative stages. Grubs fed on the roots and rootlets of young rice plants, resulted in stunting and non formation of tillers. The leaves turn yellow and develop a rusty appearance and the plants eventually die. Incidence of this pest was noticed in the month of July and August. Singh and Singh (2014) also reported the occurrence of this pest on rice crop [11].

White grub, *Holotrichia consanguinea* Blanch (Curculionidae: Coleoptera) was recorded serious polyphagous pests from nursery to maturity stages of crop in western Uttar Pradesh. The adult beetles emerged out in June–July with the onset of pre monsoon showers and defoliate preferred host trees. Grubs fed on roots of plants. The second and third instars grubs are voracious feeder and destroy the entire root. As a result the plant withered and died. Kashyap (2013) also reported the incidence of this insect from July to October on various *Kharif* crops [4].

Termite, *Odontotermes obesus* Romb (Termitidae : Isoptera) is a polyphagous social insect, also caused damage to the rice crop by feeding on the roots of the plants. The growing shoots withered and died. The damaged plants pulled out easily. The incidence of this pest was observed throughout the crop season. Such type of effect has earlier reported on different crops by Prasad and Prasad (2006), Sachan *et al.* (2006), Kashyap (2013) and Singh and Singh (2014) [4, 8, 9, 11].

Grasshoppers, *Hieroglyphus banian* Fab. and *Oxya fuscovittata* Marshall (Acrididae : Orthoptera) were found to attack basmati rice in this region. They remained active throughout the crop season. Both nymphs and adults of grasshopper fed on the leaves by making holes. In severe infestation, the leaves are completely eaten by nymphs and adults, leaving the midrib and stalk. Both these species of grasshopper are polyphagous pest and have earlier been reported by Sachan *et al.* (2006), Prasad and Prasad (2006), Kashyap (2013), Singh and Singh (2014) and Gangwar *et al.* (2015) on paddy crop [3, 4, 8, 9, 11].

Among the predators, the spiders and mirids were the most important natural enemies. Spiders and odonata recorded as general predators of rice pests. Among the odonata, damselfly population was more compared to the dragon flies. Mirids, *Cyrtorhinus lividipennis* (Reuter) was considered as important, potential and efficient predator of BPH and WBPH. Staphylinids were identified as *Paederus fuscipes* (Curtis) which is a predator on leafhoppers. Coccinellids and Cicindellids were also found on the bunds of flowering plants. The collected specimens were identified as predatory. Other important natural enemies on the insect pests of rice recorded as parasitoid which belongs to the hymenopterans orders. The recorded hymenopterans were identified parasitoids such as,

Ischnojoppalutator (Fabricius), *Xanthopimpla punctate* (Fabricius), *Xanthopimpla* sp. (larval and pupal parasitoid of leaf folder) *Charops bicolor* (Szepligeti) (Ichneumonidae) (larval parasitoid of skipper) and *Stenobracon nicevillei* (Bingham) (pupal parasitoid of yellow stem borer) and *Apanteles* sp. (Braconidae) (larval and pupal parasitoid of leaf folder) (Table- 2), These observations are in agreement with studies by Gangwar *et al.* (2015) and Kumar *et al.* (2013) [3, 6]. In the present study, predators such as spiders, dragon and damsel flies, mirid bugs, coccinellids, carabids and cicindellids were found throughout the crop growing period with little fluctuation in three methods. However, spiders, dragon fly, damsel fly and coccinellids were more during the vegetative stage of the crop, where as mirids, Staphylinids and cicindellids were more during reproductive stage of the

crop. All the recorded predators and their population are known to be directly related to their prey population. These observations are in agreement with studies by Parasappa *et al.* (2017) [7].

4. Conclusion

This study revealed a large number of insect species associated with rice crop, portends to be a potential danger to rice production. However, the large number of natural enemies species (parasitoids and predators) also recorded in rice ecosystem. All the recorded predator and parasitoids are known to be directly related to their prey population. It suggests that important biological control agents can be exploited in the management of the major insect pests of rice.

Table 1: Insect pests associated with basmati rice during *kharif* 2017 and 2018.

Order	Family	Common name	Scientific name	Damaging stage of the pest	Severity the pests
Lepidoptera	Pyralidae	Yellow stem borer	<i>Scirpophaga incertulas</i> (Walker)	Larvae	Severe
Lepidoptera	Pyralidae	Leaf folder	<i>Cnaphalocrocis medinalis</i> (Guenee)	Larvae	Severe
Lepidoptera	Noctuidae	Swarming caterpillar (Army worm)	<i>Spodoptera mauritia</i> (Boisduval)	Larvae	Low
Homoptera	Cecadellidae	Green leaf hopper	<i>Nephotettix virescens</i> (Distant)	Nymphs and adults	Low
Homoptera	Delphacidae	Brown plant hopper	<i>Nilaparvata lugens</i> (Stal.)	Nymphs and adults	Moderate
Homoptera	Delphacidae	White Backed plant hopper	<i>Sogatella furcifera</i> (Horvath)	Nymphs and adults	Low
Hetroptera	Coreidae	Rice gundhi bug	<i>Leptocorisa acuta</i> (Thumb.)	Nymphs and adults	Low
Coleoptera	Curculionidae	Rice Root Weevil	<i>Echinocnemus oryzae</i> (Marshall)	Grubs and adults	Low
Coleoptera	Curculionidae	White grub	<i>Holotrichia consanguinea</i> (Blanch)	Grubs and adults	Moderate
Isoptera	Termitidae	Termite	<i>Odontotermes obesus</i> (Romb.)	Worker	Moderate
Orthoptera	Acrididae	Kharif grass hopper	<i>Hieroglyphus banian</i> (Fab.)	Nymphs and adults	Moderate
Orthoptera	Acrididae	Grass hopper	<i>Oxya fuscovittata</i> (Marshall)	Nymphs and adults	Low

Table 2: Natural enemies associated with basmati rice during *kharif* 2017 and 2018.

Insect order	Family	Scientific name	Host
Coleoptera	Carabidae	<i>Pheropsophus</i> sp.	General predator
Coleoptera	Carabidae	<i>Ophio neaindica</i> (Thunberg)	General predator
Coleoptera	Coccinellidae	<i>Coccinella transversalis</i> (Fabricius)	General predator
Coleoptera	Coccinellidae	<i>Cheilomenes sexmaculata</i> (Fabricius)	General predator
Coleoptera	Cicindelidae	<i>Cicindela exguttata</i> (Fabricius)	Nymphs of <i>Leptocorisa oratorius</i>
Coleoptera	Staphylinidae	<i>Paederus fuscipes</i> (Curtis)	Leafhopper
Hemiptera	Miridae	<i>Cyrtorhinus lividipennis</i> (Reuter)	Nymphs and adults of <i>Nilaparvata lugens</i>
Odonata	Coenagrionidae	Cerigrion sp.	General predator
Odonata	Coenagrionidae	<i>Ischnura aurora</i> (Brauer)	General predator
Hymenoptera	Ichneumonidae	<i>Ischnojoppalutator</i> (Fabricius)	Larval and pupal parasitoid of <i>Cnaphalocrocis medinalis</i>
Hymenoptera	Ichneumonidae	<i>Xanthopimpla punctata</i> (Fabricius)	pupal parasitoid of <i>Cnaphalocrocis medinalis</i>
Hymenoptera	Ichneumonidae	<i>Charops bicolor</i> (Szepligeti)	Larval parasitoid of <i>Pelopidas methias methias</i>
Hymenoptera	Braconidae	<i>Stenobracon nicevillei</i> (Bingham)	Pupal parasitoid of <i>Scirpophaga incertulus</i>
Hymenoptera	Braconidae	<i>Apanteles</i> sp.	Larval and pupal parasitoid of <i>Cnaphalocrocis medinalis</i>
Arachnida	Lycosidae	<i>Lycosa pseudoannulata</i> (Boesenberg & Strand)	General predator

5. Acknowledgement

The authors are thankful to Head of Department of Entomology, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, and entire faculty for providing necessary facilities and laboratory for conducting the present investigation.

6. References

1. Anonymous. Food and agriculture organization, <http://faostat.fao.org>, 2017.
2. Gowda D, Gubbaiah KS. Insect pests of rice and their management in Karnataka state of India- a review. Agricultural Research. 2011; 32(1):55-62.
3. Gangwar RK, Javeria S, Yadav K, Tyagi S, Singh R. Survey and surveillance of major insect-pests of basmati rice in Western Uttar Pradesh (India). International Journal of Research in Applied, Natural and Social Science. 2015; 3(3):1-8.
4. Kashyap AK. Evaluation of some novel insecticides against yellow stem borer, *Scirpophaga incertulus* (Walker) and rice leaf folder, *Cnaphalocrocis medinalis* (Guenee) and their effect on natural enemies in basmati rice. M.Sc. (Ag.) Thesis S.V.P. Univ. Of Agric & Tech., Meerut, 2013.
5. Kumar V, Patil BV. Insect pest fauna to rice in Tungabhadra Project area of Karnataka, during Kharif season. Karnatka Journal Agricultural Sciences. 2004; 17(3):580-581.

6. Kumar D, Raghuraman M, Singh J, Waza SA, Kumar K. Occurrence of insect-pests and natural enemies in rice (*Oryza sativa* L.) agro-ecosystem. International Journal of Plant Protection. 2013; 6(2):266-270.
7. Parasappa HH, Reddy N, Neelakanth. Rice insect pests and their natural enemies complex in different rice ecosystem of Cauvery command areas of Karnataka. Journal of Entomology and Zoology Studies. 2017; 5(5):335-338.
8. Prasad R, Prasad D. Account of insect pest problem in rice ecosystem in Ranchi. Indian Journal Entomology. 2006; 68(3):240-246.
9. Sachan SK, Singh DV, Chaudhary AS. Seasonal abundance of insect pests associated with basmati rice. Annals Plant Protection Sciences. 2006; 14(1):218-273.
10. Saini UP, Sachan SK, Singh APB, Kumar K. Insect pests associated with basmati rice in western plain zone of Uttar Pradesh, India. Plant Archives. 2015; 15(2):775-777.
11. Singh BB, Singh R. Major rice insect pests in north eastern Uttar Pradesh. International Journal Life Sciences. Biotech. & Pharma. Research. 2014; 3(1):124-143.