



E-ISSN: 2320-7078

P-ISSN: 2349-6800

www.entomoljournal.com

JEZS 2020; 8(2): 1047-1053

© 2020 JEZS

Received: 13-01-2020

Accepted: 15-02-2020

L Soro

Jean Lorougnon Guédé
University (UJLoG), Daloa P.O.
Box 150, Côte d'Ivoire

S Soro

Jean Lorougnon Guédé
University (UJLoG), Daloa P.O.
Box 150, Côte d'Ivoire

NL Yeboue

Jean Lorougnon Guédé
University (UJLoG), Daloa P.O.
Box 150, Côte d'Ivoire

O Germain Ochou

National Center for Agronomic
Research (CNRA), Station of
Bouaké, Entomology laboratory,
Abidjan 13 P.O. Box 150,
Côte d'Ivoire

Corresponding Author:**L Soro**

Jean Lorougnon Guédé
University (UJLoG), Daloa P.O.
Box 150, Côte d'Ivoire

Spatial and temporal dynamics of Coccinellidae (Coleoptera) and Formicidae (Hymenoptera): Two families of beneficial insects, in cotton fields in the Tchologo region (Northern Côte d'Ivoire)

L Soro, S Soro, NL Yeboue and O Germain Ochou

Abstract

This study was conducted to analyse spatio-temporal dynamics and the impact of rainfall and temperature on two beneficial insect-families in cotton farms in Côte d'Ivoire, Formicidae and Coccinellidae. They contribute to reduce the pest populations in the cotton farms. Data were collected in four localities treated with chemical vulgarized. Sampling, of rainfall and temperature data were collected respectively by direct counting, capture with mowing net, traps, rain gauges and thermo-hygro-meters installed. We have inventoried 7675 insects including 93% for the Formicidae and 7% for the Coccinellidae. Analysis of variance, showed that there is a significant difference in abundance of Formicidae but there is no significant difference between the population of Coccinellidae. Rainfall and temperature have not affected the dynamics of beneficial insects populations. Knowledge of those beneficial insects is important to develop an integrated cotton pests control methods.

Keywords: Cotton, beneficial insects, Formicidae, Coccinellidae, Côte d'Ivoire

1. Introduction

Cotton is most fiber's crop grown in the world. Approximately, 26 million tons of cotton fiber are produced annually^[1]. China, United States, India and Pakistan produces more than 60% of cotton fiber^[2]. Worldwide, West Africa produce only 5% of international production. In Africa countries the cotton contributes considerably to the employment and income of rural populations^[3, 4].

Cotton play a strategic position in the agricultural economy of the savannah areas, because of his important role to the socio-economic development of these populations^[4, 1]. In addition, cotton cultivation has contributed to modernize the agriculture in these areas through mechanization and crop intensification. It also contribute to the improvement of the living conditions of rural populations, the organization of cooperative movement and industrial development through the construction of factories^[5, 4]. However, cotton production is faced to many constraints that deteriorate its production and causes economical drop for producers and cotton companies. In particular, the parasitism caused by insects and other arthropods is particularly important and diversified^[6]. They cause a significant drop of the yield, in a proportion of 50 to 75% without phytosanitary protection^[7-10]. According to Matthews^[11] mentioned by Castella^[12], cotton predatory insects includes no less than 1000 species, some of them are present in all production areas in the world. Several control strategies have been developed in order to maintain pest population at economically acceptable levels^[13, 9]. These include chemical control, agricultural practices and varietal control. However, although many studies have shown the importance of beneficial insects in the context of integrated protection. This method is still very limited and vulgarised in rural areas in Côte d'Ivoire, particularly in cotton growing zones. A better knowledge of the main beneficial insects present in cotton fields in Côte d'Ivoire and particularly in the Tchologo region and their role in crop protection should be a first step towards developing an integrated pests control program in a context of reducing pesticide use and environment preservation. The main objective of the present study was to evaluate the spatio-temporal dynamics and the impact of the rain and temperature on Formicidae and Coccinellidae populations. Specifically, the objective is to forecast population variations and to understand the environmental influences on their dynamics. This will allow the action of beneficial insects to be taken into account in an integrated pest management program.

2. Materials and Methods

2.1 Study sites

The work was realized in the cotton cultivation areas, (9° 35' 00" north, and 5° 11' 00" west) located in the north of Côte d'Ivoire (Figure 1). The climate is Sudanese type with two

main seasons. A dry season from November to April characterized by harmattan with peaks in December and February and a rainy season from May to October with peaks in August and September. Annual rainfall fluctuates between 1000 mm and 1400 mm (Figure 2).




Fig 1: Study sites




Fig 2: Umbro-thermal diagram

2.2 Choice of plots by sites

In the Tchologo region, four (4) cotton production areas were selected and forty (40) farmers plots were chosen, with ten (10) plots per zone. The farmer plots were distributed

proportionally over the main sowing periods in the region. Each observation plot measures a quarter (1/4) of a hectare, or (2500 m²).

2.3 Insects sampling

Capture with mowing nets and collection by hand method have been used for the sampling. According to Kumar ^[14] a significant number of samples can justify the precision of the results obtained. Insect collections were made between July and October 2018, using a mowing net and pitfall, during the period of insect activity in the cotton farms. All samples were collected at least 10 m from the field borders, in order to capture only insects living inside the plots. Adult of Coccinellidae were captured using mowing nets or by hand. Formicidae and Coccinellidae larvae were captured using plastic pitfall inserted into the ground and filled with a preservation liquid. In each plot, two traps were placed 10 m from each other. The datas were collected every week between July and October. On each observation plot, a series of 15 counts of beneficial insects was also carried out on average from 30th days after rising, at regular intervals of one week, until the 128th day of the plant cycle. Beneficial insects were counted plant by plant, on a sample of 30 plants taken in groups of 5 consecutive plants per row, following the sequential method known as the 'diagonal' method ^[15]. beneficial insects were collected by four (4) agents from the Research and Development Department of the cotton company. They have been trained for this purpose. Rainfall and temperature datas were also collected.

2.4 Data analysis

The data obtained were subjected to a one-factor analysis of variance (ANOVA 1) at the 5% threshold and ecological indices of gama and alpha diversity were determined using Past 3.2.6 software.

3. Results

3.1 Data sampling

We have inventoried 7675 insects (Table 1). The Formicidae is the most important family with 6373 individus collected, that represent 93% of the insects. The highest numbers of insects were collected in Kong (59%) and Diawala (19%) localities. Specifically, the species *Camponotus maculatus*, *Camponotus sericeus* and *Messor galla* are more abundant with respectively (28%, 26% and 24% species), followed by *Pheidole* sp (10% species). Coccinellidae are present with 1302 species, that represent 7% of all insects captured. *Cheilomenes sulphurea* is the most important species in all the cotton plots with 1044 individus collected, it represents 80% of the Coccinellidae. *Delphatus pusillus* is the least abundant with 61 individus, which is 5% of the total number of Coccinellidae. The localities of Kong (722 individus) and Korokara (277 individus) have the highest number of Coccinellidae of all species combined. Diawala (186 individus) and Kaouara (117 individus) have the lowest abundance of Coccinellidae.

Table 1: Distribution of insects inventoried by order, families and species according to localities in the Tchologo region

Order	Families	Species	Korokara	Diawala	Kaouara	Kong	Total areas
Hymenoptera	Formicidae	<i>Camponotus maculatus</i>	32	313	149	1272	1766
		<i>Camponotus acvapimensis</i>	88	8	0	199	295
		<i>Camponotus sericeus</i>	233	486	107	826	1652
		<i>Palothureus tarsatus</i>	32	0	0	18	50
		<i>Messor galla</i>	172	351	102	928	1553
		<i>Pheidole</i> sp	177	67	134	232	610
		<i>Monomorium dakarene</i>	32	19	0	355	406
		<i>Monomorium afrum</i>	41	0	0	0	41
Coleoptera	Coccinellidae	<i>Cheilomenes sulphurea</i>	216	157	117	554	1044
		<i>Cheilomenes propinqua vicini</i>	38	19	0	140	197
		<i>Delphatus pusillus</i>	23	10	0	28	61
Total			1084	1430	609	4552	7675
Percentage of Formicidae			74%	94%	68%	84%	93%
Percentage of Coccinellidae			26%	6%	32%	16%	7%

3.2 Spatial distribution of Formicidae and Coccinellidae

Presence of Formicidae was observed in all the plots. The highest abundance of insects was recorded in Kong, with dominance of the species *Camponotus maculatus* (1272 individus), *Camponotus sericeus* (826 individus), *Messor galla* (928 individus) followed by *Pheidole* sp and *Monomorium dakarene* respectively with 232, 355 individus. All the species were observed in all sites with mixed numbers. However, at Kaouara the lowest abundance of insects was recorded with total absence of *Camponotus acvapimensis*, *Palothureus tarsatus*, *Monomorium dakarene* and *Monomorium afrum* (Figure 3). The specific richness (S) of Formicidae is important in all the localities. The The highest number of species was recorded in Korokara (8 species), followed by Kong (7 species). Shannon wiener (H') and Pielou (E) index values trend in the same direction as the

specific richness (Table 2). Analysis of variance show that there is significant difference between abundance of species captured in the localities (df = 31, F = 4.08 and P = 0.015). Coccinillidae was captured in all the localities. *Cheilomenes sulphurea* is the most abundant specie with mixed numbers. *Delphatus pusillus* is rare in the plots with an absence in Kaouara locality. Korokara, Diawala and Kong sites, recorded the greatest specific richness with 3 species while in Kaouara locality only one species was captured (Figure 4). The low values of Pielou equitability index vary between 0.48 and 0.61 (Table 3). Its show that the distribution of Coccinellidae in the different localities is not equal between them. Analysis of variance show that there is no significant difference between abundance of species captured in the localities (df = 11, F = 0.99 and P = 0.44).




Fig 3: Spatial distribution of Formicidae




Fig 4: Spatial distribution of Coccinellidae

Table 2: Ecological index of Formicidae

Localities	Specific Richness (S)	Individus (N)	Shannon Weaner (H')	Pielou equitability (E)
Korokara	8	807	1.798	0.864
Diawala	6	1344	1.292	0.721
Kaouara	4	492	1.374	0.991
Kong	7	3830	1.610	0.827

Table 3: Ecological index of Coccinellidae

Localities	Specific Richness (S)	Individus (N)	Shannon Weaner (H')	Pielou equitability (E)
Korokara	3	277	0.673	0.612
Diawala	3	186	0.533	0.485
Kaouara	1	117	0	
Kong	3	722	0.643	0.589

3.3 Weekly fluctuations of Formicidae and Coccinellidae

The most important peak of Formicidae captures started the last week of July (143 individus of Formicidae) in Korokara (Figure 5a). Formicidae abundance dropped in October compared to the captures of Coccinellidae which were observed during the period of the study, from July to October. Weekly fluctuations of the two beneficial insects families observed in Diawala locality (Figure 5b) show that the number of insects observed varies according to the weeks. The most important peak of Formicidae were reached during the second week of August, 2018. Indeed, this important

population would be due to the infestation of cotton plots by aphids in August. In Kaouara plots, Formicidae populations were present during the study period, Coccinellidae were observed from July to the end of September with two peaks in August, the most important one (36 individus) during the 3rd week of August 2018 (Figure 5c). In Kong cotton plots, Formicidae and Coccinellidae were observed from the beginning of the inventories. The first peak of Formicidae was observed as early as the third week of capture (602 individus). The number of Coccinellidae was relatively constant from the beginning to the end of the observations in October 2018.




Fig 5: Weekly fluctuations of Formicidae and Coccinellidae

3.4 Influence of Rainfall on Population Dynamics

August and September 2018 were more watered. Overlay graphs of the temporal distribution of Formicidae and Coccinellidae and the evolution of rainfall shows that peaks of insects populations are independent of rainfall in all localities (Figure 6). The dynamics of the Formicidae and Coccinellidae are not influenced by rainfall.

3.5 Influence of temperature on population dynamics

Temperature data were relatively constant over the study period. The temperature fluctuated between 24 °C and 26 °C. Overlaying the graphs (Figure 6) shows that there is no coincidence between the temperature variations recorded during the surveys and the fluctuations of the Formicidae and Coccinellidae populations.




Fig 6: Influence of rainfall and temperature on the dynamics of Formicidae and Coccinellidae

4. Discussion

This work, conducted in 2018, made it possible to collect an important but probably non-exhaustive population of beneficial insects of Coleoptera and Hymenoptera from cotton plots in the Tchologo region. These are: *Camponotus* sp, *Pheidole* sp and *Messor* sp for Formicidae and *Cheilomenes* sp and *Delphatus* sp Coccinellidae. These results are similar to those obtained by Sigrist *et al* (1994) [16] in a study of the non-intentional effects of pesticides on beneficial insects of cotton in Chad. The work done by Galva (1993) [17] had also noted the presence of several beneficial insects in the cotton fields in northern Cameroon, including several species of Formicidae and Coccinellidae. Also, according to Hamiti and Bouchaala (2013) [18], the role of biocontrol crop species is important. They have the advantage to be non-toxic. The study of the spatio-temporal dynamics of the two families of the beneficial insects of cotton is by alternating high and low population densities according to time and space. Density fluctuations of insects populations observed during the study has been done according to the areas and the species. Thus, peaks of Formicidae and Coccinellidae were observed between August and September in all localities. Tuo's work in 2013 [19] showed that the peaks of palm entomofauna species vary according to the study site. In order to appreciate the influence of climatic factors on the two families of cotton beneficial insects, the variation of rainfall and temperature graphs in relation with time were compared for each family. This analysis showed that climatic factors did not significantly influence insect dynamics in the cotton plots. These results are confirmed by Mariau *et al.* (1991) [20] and Tuo (2013) [19] who indicated that there is no significant

relationship between the emergence of insects in general and climatology. The beneficial insects studied were observed throughout the cotton cycle, which could be explained by the fact that during the development cycle the climate influences the development of the plants that serve as a biotope, Chararas, (1959) [21] for the insects. According to Koné *et al* 2017 [22], the parasite distribution currently observed in the cotton production areas could be due to high rainfall intensity.

5. Conclusion

From the present investigation it can be concluded that Formicidae and Coccinellidae are among the most important beneficial insects presents in cotton farms. After the investigation in 2018, 1798 specimens have been recorded. The spatial distribution analysis show that the Formicidae and Coccinellidae are present in all the study localities during the cycle of the cotton plants and then the climatic factors don't influence their abundance. It would therefore be important to take into consideration the action of beneficial insects in the development of cotton pest control programs in Côte d'Ivoire.

6. Acknowledgement

We would like to thank the staff of Research and Development Department of SECO-SA for the collection of insects and Professor Yéo Kolo and his team of Lamto Ecological Station for the identification of specimens.

7. References

1. Aïwa A. L'impact de la culture du coton sur le développement socio-économique : étude de cas de la région de Korhogo, au nord de la cote d'ivoire. European

- Scientific Journal. 2015; 11(31):1857-7881
2. Fabio B, Jean-Luc H, Zagbaï HS, Lebailly P. Le coton dans le monde, place du coton africain et principaux enjeux. *Biotechnology, Agronomy, Society and Environment*. 2006; 10(4):271-280.
 3. Matthes A, Van Den AE, Chougourou D, Midingoyi J. S. Le coton au Bénin : Compétitivité et durabilité de cinq systèmes culturaux cotonniers dans le cadre de la filière, BMZ-GTZ/MAEP, Cotonou. 2005, 201
 4. Soro L. Dynamique cotonnière en Côte d'Ivoire et analyse des impacts socio-économiques : Cas de la SECO. Mémoire de Master professionnel du groupe IFPG-ISFPT, Négocier des matières premières agricoles. 2014, 94
 5. Zagbaï SH. Evaluation de l'impact du coton sur le développement. Cas de la région de Korhogo au nord de la Côte d'Ivoire. Thèse de doctorat. Faculté Universitaire des Sciences Agronomiques de Gembloux, Unité d'Économie et Développement Rural. 2005, 243
 6. Vaissaire M, Cauquil J, Silvie P. Protection phytosanitaire du cotonnier : Méthodes et moyens de lutte intégrée contre les ravageurs. *Agriculture et Développement*. 1995; (8):4-23.
 7. Ochou OG. Lutte contre les maladies et arthropodes du cotonnier. Note technique IDESSA-DCI / FC /PDC. 1998, 46
 8. Rafalimanana HJ. Evaluation des effets d'insecticides sur deux types d'Hyménoptères auxiliaires des cultures, l'abeille domestique (*Apis mellifera* L.) et des parasitoïdes de pucerons: études de terrain à Madagascar et de laboratoire en France. Thèse de Doctorat de De l'institut national agronomique paris- grignon. 2003 ; 206
 9. Badiane D., Gueye MT., Cole EV., Faye O., Gestion intégrée des principaux ravageurs du cotonnier au Sénégal et en Afrique occidentale. *International Journal of Biological and Chemical Sciences*. 2015; 9(5):2654-2667
 10. Sarr M, Badiane D, Sane B. Evaluation de l'efficacité de nouveaux programmes de protection phytosanitaire contre les principaux ravageurs du cotonnier *Gossypium hirsutum* L. au Sénégal. *International Journal of Biological and Chemical Sciences*. 2016; 10(5):2163-2174.
 11. Matthews GA. Cotton insect pests and their management. Harlow, Essex, UK : Longman Scientific & Technical ; New York : Wiley, 1989; 633(5):19
 12. Castella JC. Stratégies de lutte contre les insectes ravageurs dans les systèmes de culture cotonniers en Thaïlande: Logiques actuelles et propositions pour une gestion durable. Thèse de doctorat de l'Institut National Agronomique Paris-Grignon, Sciences Agronomiques, 1995; 152 (4):339
 13. FAO. Gestion intégrée de la production et des déprédateurs du coton, guide du facilitateur pour les champs écoles des producteurs, 2014; 208280(4):89
 14. Kumar R. La Lutte Contre les Insectes Ravageurs ; la Situation de l'Agriculture Africaine. Paris, France. 1991, 310
 15. Nibouche S, Beyo J, Goze E. Mise au point de plans d'échantillonnage pour la protection sur seuil contre les chenilles de la capsule du cotonnier. Colloque Savanes africaines : des espaces en mutation, des acteurs face à de nouveaux défis, Garoua, Cameroun. hal-00140842. 2003, 5
 16. Sigrist JC, Martin T, Renou A. Effets non intentionnels des pesticides sur l'entomofaune utile des cotonniers ; CIRAD, unité de recherche entomologie appliquée. 1994, 9
 17. Galva P. Contribution à l'étude des auxiliaires naturels en culture cotonnière paysanne au Nord Cameroun. Mémoire de fin d'études, CNEARC Montpellier, France, 1993, 76
 18. Hamiti Y, Bouchaala H. Inventaire des coccinelles prédatrices pouvant être utilisées dans un cadre de lutte biologique dans la région de Ouargla, Mémoire de master académique, Département des Sciences Agronomiques, Sciences de la Nature et de la Vie. 2013, 94
 19. Tuo Y. Etat de l'entomofaune des inflorescences du palmier à huile en Côte d'Ivoire : Cas de la station de la Mé. Thèse de doctorat de l'Université Felix Houphouët Boigny, entomologie agricole. 2013, 204
 20. Mariau D, Houssou M, Lecoustre R, Ndigui B. Insectes pollinisateurs du palmier à huile et taux de nouaison en Afrique de l'ouest. Oléagineux. Mémoire de synthèse des activités scientifiques présenté en vue de l'obtention de l'habilitation à diriger des recherches. Université des Sciences et Technologies de Lille. 1991; 46(2):43-49 Chararas. C. Influence des conditions climatiques sur l'évolution des Scolytides, *Compte Rendu de plusieurs années d'expérimentation dans les forêts de Haute-Savoie*. 1958, 33
 21. Koné PWE, Ochou GEC, Didi GJR, Dekoula SC, Kouakou M, Bini KKN. *et al.* Evolution spatiale et temporelle des dégâts du jasside *Jacobiella facialis* Jacobi, 1912 (Cicadellidae) en comparaison avec la distribution de la pluviométrie au cours des vingt dernières années dans les zones de culture cotonnière en Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*. 2017; 11(3):1190-1201.