



E-ISSN: 2320-7078

P-ISSN: 2349-6800

www.entomoljournal.com

JEZS 2020; 8(2): 450-454

© 2020 JEZS

Received: 12-01-2020

Accepted: 14-02-2020

Usha

Assistant Professor, Department of Entomology, College of Agriculture, Rani Lakshmi Bai Central Agricultural University, Jhansi, Uttar Pradesh, India

Vijay Kumar Mishra

Teaching cum Research Associates, College of Agriculture, Rani Lakshmi Bai Central Agricultural University, Jhansi, Uttar Pradesh, India

Maimom Soniya Devi

Teaching cum Research Associates, College of Agriculture, Rani Lakshmi Bai Central Agricultural University, Jhansi, Uttar Pradesh, India

Corresponding Author:**Usha**

Assistant Professor, Department of Entomology, College of Agriculture, Rani Lakshmi Bai Central Agricultural University, Jhansi, Uttar Pradesh, India

Effect of environmental factors on the foraging activities of major bee pollinators

Usha, Vijay Kumar Mishra and Maimom Soniya Devi

Abstract

Environment plays an important role in pollinator's activity. Flight activity in *Apis mellifera* was significantly influenced by temperature, relative humidity and sunshine. Maximum temperature exerted a significant positive correlation with the foraging activity of *Apis dorsata*, *Apis mellifera* and *Apis cerana indica* which indicate that maximum temperature fell within the optimal range (21.0 to 33.5 °C) for their activity. However, minimum temperature exerted significant positive correlation with the activity of only *A. dorsata* and *A. c. indica*. It reveals that the *A. dorsata* and *A. c. indica* are able to forage with optimal range of maximum and minimum temperature. It was also observed that maximum temperature exerted significant negative correlation with the foraging activity of syrphids and non apis bees, means their activity is decreased with increasing temperature. Although, rain fall exerted significant negative correlation with all foragers showed their activity is decreased during rainy day.

Keywords: *A. dorsata*, *A. mellifera*, *A. c. indica*, syrphids and non apis bees etc

Introduction

Insect pollinators like syrphid flies, honey bees and non apis bees have proved as boon for efficient and effective pollination in most of the crops. They are affected directly or indirectly by the environment around them. Among different aspects the environment plays an important role in pollinator's activity. Flight activity in *A. mellifera* was significantly influenced by temperature, relative humidity and sunshine. Reduced activity of pollinators due to low temperature during flowering contributed to the formation of 5.44 percent parthenocarpic seedless fruits in mango (Thimmappaiah and Singh, 1983) [22].

The low light intensity and low temperature probably discouraged foraging (Tanda, 1984) [21]. Khader (1989) [11] also observed in mango fruit setting of a cluster of abnormal seedless at the tip of the panicle called "JHUMKA" in Lucknow area. Most of these fruits contained aborted and shriveled embryos and dropped in less than two weeks. Failure of pollination due to adverse climatic conditions in February-March was considered as a major factor responsible for this. Pande *et al.* (1991) [16] noticed that the foraging activities of honeybees were maximum at higher atmospheric temperature and low relative humidity.

Abrol (1992) [1] and Corbet *et al.* (1993) [5] indicated that several environmental factors *viz.*, temperature, humidity, light, solar radiation, time of the day and nectar flow regulate flight activity of bees. Initiation of flight was influenced by dual threshold of temperature and light intensity and cessation was controlled by reduction of light intensity for bee species. Bee visitation was more intensive during late morning than early afternoon. Strength of wind was the most important factor in determining honeybee visitation. The number of pollinators/panicle/minute increased as the average temperature increased to 19.5° C but decreased with further increase in temperature and with rains (Singh *et al.*, 1997) [17]. Singh and Devi (1998) [18] worked out a positive correlation of initiation and cessation of foraging activity with temperature whereas, relative humidity had positive correlation with cessation of foraging in *A. cerana*. According to Abrol (1998) [2] the activity of both bees species, *A. cerana* and *A. mellifera* was negatively correlated with relative humidity and positively with light intensity, solar radiation, nectar sugar concentration and temperature. Ambient temperature and light intensity affected flight activity positively and relative humidity negatively. The meteorological factors were responsible for 33.5 per cent of variation in number of bees leaving the hive and 27 per cent of the variation in bees returning to hive in *A. mellifera*. Abrol (2000) [3] observed that the diurnal activity showed significant negative correlation with temperature and significant positive correlation with humidity.

Among different aspects, the environment plays very important role in bee foraging. Flight activity in case of *A. mellifera* was significantly influenced by temperature, relative humidity and sunshine. Foraging was also affected by wind speed, which varied from 0.2 to 22.89 km/h than by air temperature and relative humidity.

The main abiotic factors that affect the flight of bees are temperature, light intensity, relative air humidity, and wind speed (Kleinert-Giovannini 1982^[12]; Sommeijer *et al.* 1983^[19]; Corbet *et al.* 1993a^[6]; Hilário *et al.* 2000^[9]; Vicens and Bosch 2000^[20]; Kleinert *et al.* 2009). Extreme temperatures, either too hot or too cold, restrict flight activity (Hilário *et al.* 2000)^[9]. Flying hymenopterans dehydrate or their body temperature increases to near lethal levels when the relative humidity is too low or under conditions of very high temperature, respectively (Coelho and Ross 1996;^[4] Kasper *et al.* 2008)^[10]. Light intensity acts especially on helping visual navigation of bees, from orientation to the search of floral resources (Hilário *et al.* 2000)^[9]. At low ambient temperatures, the light intensity is also important for maintaining the body heat of the bees (Kovac *et al.* 2009^[13]; Kovac and Stabentheiner 2011^[15]; Hartfelder *et al.* 2013)^[8]. Very strong winds affect the flight activity of bees, preventing them from foraging under such conditions (Sommeijer *et al.* 1983)^[19]. In turn, the combined effect of environmental factors determines the insects' operative temperature (Kovac *et al.* 2010^[14]; Hartfelder *et al.* 2013)^[8] which is achieved from the transfer of heat by conduction, convection and radiation between the body of the insect and the environment (Dzialowski, 2005)^[7]. In an environmental temperature of 15.1 °C and solar radiation of 684 W m⁻² Kovac *et al.* (2010)^[14] recorded a mean thoracic temperature of 24.4 °C in two dead honeybee workers. Each bee species depends on specific climatic conditions, which set the boundaries for the foraging flight (Corbet *et al.* 1993a)^[6].

Keeping in view the importance of insect pollinators, a comprehensive study was undertaken to determine impact of abiotic factors on the foraging activity of the major bee pollinators.

Material and Methods

The visitation of insect pollinators were observed visually during the blooming period to record the influence of abiotic factors on the forager's activity. The data on Temperature, Relative Humidity and Rain fall were obtained from the Agrometeorology Department were correlated with the insect pollinators activity.

Results and Discussion

Data recorded on the abundance of different insect pollinators/visitors was correlated with weather parameters such as Maximum and Minimum temperature, Relative humidity and total rainfall and result have been presented in the table 1 – 4 and Fig 2 & 3. It was apparent that when all the weather factors acted together, exerted their influence differently on different foragers during the year 2018 and 2019. The multiple correlation coefficient (r) against maximum temperature (21.0-33.5 °C), minimum temperature (7.9-20.0°C), % relative humidity I hour (73-91%), II hour (22-60%) and total rainfall (00.0-9.2 mm.) was computed to be-for -0.064*, -0.055, 0.823, -0.040 and -0.442*, respectively for Syrphid flies, 0.239*, 0.190*, 0.513, -0.271 and -0.265* for *A. dorsata*, 0.240*, 0.282*, 0.468, -0.410 and -0.153* for *A. mellifera*, 0.035*, 0.024*, 0.649, -0.025 and -

0.364* for *A. cerana indica* and -0.456*, -0.386, 0.977, 0.311 and -0.574* for Non apis bees (Table 1) during the year 2018. Observations revealed that maximum temperature exerted a significant positive correlation with the foraging activity of *A. dorsata*, *A. mellifera* and *A. c. indica* which indicate that maximum temperature fell within the optimal range (21.0 to 33.5°C) for their activity. However, minimum temperature exerted significant positive correlation with the activity of only *A. dorsata* and *A. c. indica*. It reveals that the *A. dorsata* and *A. c. indica* are able to forage with optimal range of maximum and minimum temperature. It was also observed that maximum temperature exerted significant negative correlation with the foraging activity of syrphids and non apis bees, means their activity is decreased with increasing temperature. Although, rain fall exerted significant negative correlation with all foragers showed their activity is decreased during rainy day. While, relative humidity showed no significant effect for any foragers.

The combined effects of all the parameters had been summarized in Table 2. It was apparent that when all the weather factors acted together, exerted their influence differently on different foragers. The coefficient of determination (R²) was computed to be 0.004, 0.057, 0.058, 0.001 and 0.208 against Syrphid flies, *A. dorsata*, *A. mellifera*, *A. c. indica* and Non apis bees controlling 0.40, 5.70, 5.80, 0.10 and 20.80 per cent visitors foraging activity alone, respectively. Similarly, from the Table 3 the impact of weather variables significantly affected the visitation intensities of the honeybee, *A. mellifera*, *A. dorsata*, *A. cerana indica*, Non- Apis Bees and Syrphid flies during the blooming period of the mango in the year 2019. The multiple correlation coefficient (r) against maximum temperature (27.5-35.7°C), minimum temperature (11.0-18.7°C), % relative humidity I hour (72-90%), II hour (30-49%) and total rainfall (00.0-007.4mm) was computed to be:-0.288*, -0.477, 0.711, 0.092 and -0.412* for Syrphid flies, 0.326*, 0.151*, 0.238, -0.131 and -0.451* for *A. dorsata*, -0.057*, -0.197, 0.591, -0.149 and -0.304* for *A. mellifera*, 0.195*, 0.0004*, 0.328, -0.114 and -0.466* for *A. cerana indica*, -0.074*, -0.243, 0.596, 0.080 and -0.362* for Non apis bees. Results revealed that maximum temperature exerted a significant positive correlation with the foraging activity of *A. dorsata* and *A. c. indica* which indicate that maximum temperature fell within the optimal range (27.5 to 35.7°C) for their activity. However, minimum temperature exerted significant positive correlation with the activity of only *A. dorsata* and *A. c. indica*. It reveals that the *A. dorsata* and *A. c. indica* are able to forage with optimal range of maximum and minimum temperature. It was also observed that maximum temperature exerted significant negative correlation with the foraging activity of syrphids, *A. mellifera* and non apis bees, means their activity is decreased with increasing temperature. Although, rain fall exerted significant negative correlation with all foragers showed their activity is decreased during rainy day. While, relative humidity showed no significant effect for any foragers. Similarly Pande *et al.* (1991)^[16] noticed that the foraging activities of honeybees were maximum at higher atmosphere temperature and low relative humidity. The above findings were supported by the experimental findings of Singh *et al.* (1997)^[17] who reported that the number of pollinators/panicle/minute increased as the average temperature increased to 19.5° C but decreased with further increase in temperature and with rains.

The combined effects of all the parameters had been

summarized in Table 4. It was apparent that when all the weather factors acted together, exerted their influence differently on different foragers. The coefficient of determination (R²) was computed to be 0.083, 0.106, 0.003,

0.038 and 0.005 against Syrphid flies, *Apis dorsata*, *Apis mellifera*, *Apis cerana indica* and Non *Apis* bees controlling 8.30, 10.60, 0.30, 3.80 and 0.50 per cent visitors foraging activity alone, respectively.

Table 1: Correlation co-efficient matrix (r) between insect pollinators on mango and weather parameters during the year 2018

Weather parameter	Insect pollinators (r)				
	Y ₁	Y ₂	Y ₃	Y ₄	Y ₅
X ₁	-0.064*	0.239*	0.240*	0.035*	-0.456*
X ₂	-0.055	0.190*	0.282*	0.024*	-0.386
X ₃	0.823	0.513	0.468	0.649	0.977
X ₄	-0.040	-0.271	-0.410	-0.025	0.311
X ₅	-0.442*	-0.265*	-0.153*	-0.364*	-0.574*

X₁ - Maximum Temperature °C X₂ - Minimum Temperature °C X₃ - % Relative Humidity I hr.
 X₄ - % Relative Humidity II hr. X₅ - Total rainfall (mm)
 Y₁ - Syrphids flies Y₂ - *Apis dorsata* Y₃ - *Apis mellifera*
 Y₄ - *Apis cerana indica* Y₅ - Non *Apis* bees

Table 2: Multiple regression equation between dependent and independent variables during the year 2018

Insect pollinators visited	Multiple equation	Coefficient of determination (R ²)
Y ₁	29.585 - 0.039X ₁ - 0.031X ₂ + 0.939X ₃ - 0.686X ₄ - 0.222X ₅	0.004*
Y ₂	28.354 + 0.371X ₁ + 0.275X ₂ + 1.496X ₃ - 1.174X ₄ - 0.341X ₅	0.057*
Y ₃	27.618 + 0.134X ₁ + 0.146X ₂ + 0.492X ₃ - 0.638X ₄ - 0.071X ₅	0.058
Y ₄	29.145+0.147X ₁ +0.095X ₂ +4.992X ₃ -0.289X ₄ -1.236X ₅	0.001
Y ₅	33.374-1.646X ₁ -1.297X ₂ +6.619X ₃ +3.116X ₄ -1.713X ₅	0.208*

X₁ - Maximum Temperature °C X₂ - Minimum Temperature °C X₃ - % Relative Humidity I hr.
 X₄ - % Relative Humidity II hr. X₅ - Total rainfall (mm)
 Y₁ - Syrphid flies Y₂ - *Apis dorsata* Y₃ - *Apis mellifera*
 Y₄ - *Apis cerana indica* Y₅ - Non *Apis* bees

Table 3: Correlation co-efficient matrix (r) between insect pollinators on mango and weather parameters during the year 2019

Weather parameter	Insect pollinators (r)				
	Y ₁	Y ₂	Y ₃	Y ₄	Y ₅
X ₁	-0.288*	0.326*	-0.057*	0.195*	-0.074*
X ₂	-0.477	0.151*	-0.197	0.0004*	-0.243
X ₃	0.711	0.238	0.591	0.328	0.596
X ₄	0.092	-0.131	0.149	-0.114	0.080
X ₅	-0.412*	-0.451*	-0.304*	-0.466*	-0.362*

X₁ - Maximum Temperature °C X₂ - Minimum Temperature °C X₃ - % Relative Humidity I hr.
 X₄ - % Relative Humidity II hr. X₅ - Total rainfall (mm)
 Y₁ - Syrphid flies Y₂ - *Apis dorsata* Y₃ - *Apis mellifera*
 Y₄ - *Apis cerana indica* Y₅ - Non *Apis* bees

Table 4: Multiple regression equation between dependent and independent variables during the year 2019

Insect pollinators visited	Multiple equation	Coefficient of determination (R ²)
Y ₁	33.302-0.097X ₁ -0.161X ₂ +0.566X ₃ +0.070X ₄ -0.139X ₅	0.083*
Y ₂	31.343+0.563X ₁ +0.262X ₂ +0.973X ₃ -0.518X ₄ -0.781X ₅	0.106*
Y ₃	32.535-0.027X ₁ -0.094X ₂ +0.665X ₃ +0.162X ₄ -0.145X ₅	0.003
Y ₄	31.870+0.772X ₁ -0.001X ₂ +3.061X ₃ -1.027X ₄ -1.846X ₅	0.038
Y ₅	32.554-0.184X ₁ -0.601X ₂ +3.478X ₃ +0.450X ₄ -0.897X ₅	0.005*

X₁ - Maximum Temperature °C X₂ - Minimum Temperature °C X₃ - % Relative Humidity I hr.
 X₄ - % Relative Humidity II hr. X₅ - Total rainfall (mm)
 Y₁ - Syrphid flies Y₂ - *Apis dorsata* Y₃ - *Apis mellifera*
 Y₄ - *Apis cerana indica* Y₅ - Non *Apis* bees




Fig 2: Influence of weather parameters on foraging behaviour of insect pollinators during the year 2018




Fig 3: Influence of weather parameters on foraging behaviour of insect pollinators during the year 2019

Conclusion

Observations revealed that maximum temperature exerted a significant positive correlation with the foraging activity of *A. dorsata*, *A. mellifera* and *A. c. indica* which indicate that maximum temperature fell within the optimal range (21.0 to 33.5°C) for their activity. However, minimum temperature exerted significant positive correlation with the activity of only *A. dorsata* and *A. c. indica*. It reveals that the *A. dorsata* and *A. c. indica* are able to forage with optimal range of maximum and minimum temperature. It was also observed that maximum temperature exerted significant negative correlation with the foraging activity of syrphids and non APIs bees, means their activity is decreased with increasing temperature. Although, rain fall exerted significant negative correlation with all foragers showed their activity is decreased during rainy day. While, relative humidity showed no significant effect for any foragers.

References

- Abrol DP. Foraging in honeybees *Apis cerana indica* F. and *A. dorsata* F. (Hymenoptera: Apidae) activity and weather conditions. *Indian Inst. Sci.*, 1992; 72(5):395-401.
- Abrol DP. Environmental factors influencing flight activity in honey bees, *A. cerana* Fab. and *A. mellifera* L. (Hymenoptera: Apidae). *Indian Bee Journal*, 1998; 60(2):71-75.
- Abrol DP. Insect pollinators of kiwifruit (*Actinida deliciosa*). *Indian Bee Journal*. 2000; 62(3-4):55-57.
- Coelho JR, Ross AJ. Body Temperature and Thermoregulation in two Species of Yellowjackets *Vespa Germanica* and *V. Maculifrons*. *J Comp Physiol B Biochem Syst Environ Physiol*. 1996; 166:68-76.
- Corbet Sarah A, Williams Ingrid H, Osborne Juliet L. Bees and the pollination of crops and wild flower in the European community. *Bee World*, 1993, 47-59.
- Corbet SA, Fussel M, Ake R, Fraser A, Gunson C, Savage A *et al*. Temperature and Pollination Activity of Social Bees. *Ecol Entomol*. 1993a; 18:17-30.
- Dzialowski EM. Use of Operative Temperature and Standard Operative Temperature Models in Thermal Biology. *J Therm Biol*. 2005; 30:317-334.
- Hartfelder K, Bitondi MMG, Brent CS, Guidugli-Lazzarini KR, Simões ZLP, Stabentheiner A *et al*. Standard methods for physiology and biochemistry research in *Apis mellifera*. In: Dietemann V, Ellis JD, Neumann P (eds) *The coloss beebook*, vol. I: standard methods for *Apis mellifera* research. *J Apic Res*. 2013; 52:26-31.
- Hilário SD, Imperatriz-Fonseca VL, Kleinert AMP. Flight Activity and Colony Strength in the Stingless bee *Melipona Bicolor Bicolor* (Apidae, Meliponinae). *Rev Bras Biol*. 2000; 60:299-306.
- Kasper ML, Reeson AF, Mackay DA, Austin AD. Environmental Factors Influencing Daily Foraging Activity of *Vespa Germanica* (Hymenoptera, Vespidae) in Mediterranean Australia. *Insectes Soc*. 2008; 55:288-295.
- Khader S. A short note on clustering in mango a new disorder. *Haryana. J Hort. Sci*. 1989; 18:233-234.
- Kleinert-Giovannini A. The Influence of Climatic Factors on Flight Activity of *Plebeia Emerina* Friese (Hymenoptera, Apidae, Meliponinae) in winter. *Rev Bras Entomol*. 1982; 26:1-13.

13. Kovac H, Stabentheiner A, Schmaranzer S. Thermoregulation of Water Foraging Wasps (*Vespula Vulgaris* and *Polistes Dominulus*). *J Insect Physiol.* 2009; 55:959-966.
14. Kovac H, Stabentheiner A, Schmaranzer S. Thermoregulation of Water Foraging Honeybees: Balancing of Endothermic Activity with Radiative Heat Gain and Functional Requirements. *J Insect Physiol.* 2010; 56:1834-1845.
15. Kovac H, Stabentheiner A. Thermoregulation of Foraging Honeybees on Flowering Plants: Seasonal Variability and Influence of Radiative Heat Gain. *Ecol Entomol.* 2011; 36:686-699.
16. Pande P, Panda B, Sontake BK. Foraging behaviour of honey bees on sunflower (*Helianthus annuus* L.) sown on different dates. *Indian Bee J.* 1991; 53(1-4):44-49.
17. Singh G, Lavi V, Degani C, Gazit Lahav S, Pesisau E, Prusky D *et al.* Pollination, pollinators and fruit setting in mango. *Proceedings 5th International Symposium on mango*, Tel Aviv, Israel, 1997, 16-123.
18. Sing MP, Devi CS. Effect of environmental factors on the activities of *Apis cerana* F. in rape seed pollination. *Insect Environ.* 1998; 4(2):50.
19. Sommeijer MJ, Derooy GA, Punt W, Bruijn LLM. A Comparative Study of Foraging Behavior and Pollen Resource of Various Stingless Bees (Hym. Meliponinae) and Honey Bees (Hym. Apinae) in Trinidad, West-Indies. *Apidologie.* 1983; 14:205-224.
20. Vicens N, Bosch J. Weather-Dependent Pollinator Activity in an Apple Orchard, With Special Reference to *Osmia Cornuta* and *Apis Mellifera* (Hymenoptera: Megachilidae and Apidae). *Environ Entomol.* 2000; 29:413-420.
21. Tanda AS. Forging behaviour of three species of *Apis* on raya relation to the sugar concentration in its nectar. *Indian Bee Journal*, 1984; 46:5-6.
22. Thimmappaiah, Harmail Singh. Natural parthenocarpy in Dashehari mango. *Indian J Hort.* 1983; 40(3/4):195-198.