



E-ISSN: 2320-7078

P-ISSN: 2349-6800

www.entomoljournal.com

JEZS 2020; 8(5): 1598-1600

© 2020 JEZS

Received: 24-06-2020

Accepted: 22-08-2020

AY Munj

Jr. Entomologist, Regional Fruit
Research Station, Vengurle
Maharashtra, India

PV Reddy

Principal Scientist, Indian
Institute of Horticultural
Research, Bengaluru,
Karnataka, India

Gundappa

Scientist, Central Institute for
Subtropical Horticulture,
Lucknow, Uttar Pradesh, India

JK Bana

Jr. Entomologist, Agricultural
Experiment Station, Paria,
Gujarat, India

S Irulandi

Jr. Entomologist, Horticulture
College and Research Institute,
Periyakulam, Tamil Nadu, India

Management of mango hopper using entomopathogens

AY Munj, PV Reddy, Gundappa, JK Bana and S Irulandi

Abstract

Mango hopper is an important pest of mango in India, causing severe yield losses. Many insecticides have been recommended so far for the management of mango hopper. These inorganic synthetic insecticides have many side effects including resurgence, resistance, residue, etc. Therefore, there is a need to recommend eco-friendly control measures for the management of mango hopper. Experiments were conducted at different mango research centers in different states of India for management of mango hopper with different entomopathogens during 2015-16 to 2017-18. The results revealed that the treatment of five sprays of *Metarrhizium anisopliae* (oil formulation @ 0.5 ml/l.) supplied by IIHR, Bengaluru was the most effective at Bengaluru and Periyakulam, whereas, the treatment of five sprays of *Verticillium lecanii* (Commercial product @ 5 gm/l.) was the most effective at Lucknow, Paria and Vengurle.

Keywords: Mango hopper, management, entomopathogens, *Metarrhizium anisopliae*, *Verticillium lecanii*

Introduction

Mango, *Mangifera indica* L. is an important fruit crop grown in India. It is found to be infested by more than 50 insect pests^[1]. Mango hopper is one of the most serious pests responsible for about 60 percent losses in fruit yield^[2]. Three species of mango hoppers viz. *Amritodus atkinsoni*, *Idioscopus clypealis* and *I. nitidulus* are commonly found all over India^[3].

The female hoppers lay eggs in the midrib on the under surface of tender leaves and on the tender panicles at the time of panicle initiation. The incubation period is 3 to 5 days and nymphal period is 10 to 15 days^[4]. The hopper incidence is severe during November to March in different parts of India^[5]. The adult hoppers as well as nymphs damage all the tender part of mango viz., tender foliage, inflorescence and fruits. They suck cell sap from these plant parts, as a result there is twisting of tender leaves, blackening of inflorescence and shading of flowers. Also, the yield is badly affected. In addition, hoppers excrete honey dew like substance on which black sooty mould (*Capnodium mangiferae*) grows which interferes with the photosynthetic activity of leaves and reduces the market value of fruits^[6].

Many inorganic insecticides have been recommended so far, for management of mango hopper throughout India^[7-13]. However, harmful chemical insecticides used for the management of mango hopper created many problems like resistance, resurgence, residue among others.^[14] Therefore, there is a need for eco-friendly practices for the management of mango hopper. The present study was conducted using different entomopathogens under the All India Co-ordinated Research Project on Fruits at five research institutes viz., IIHR-Bengaluru (Karnataka), CISH-Lucknow (Uttar Pradesh), AES-Paria (Gujarat), CHRI-Periyakulam (Tamil Nadu) and RFRS-Vengurle (Maharashtra).

Materials and Methods

The management trials were conducted at the five research institutes in 2015-16, 2016-17 and 2017-18. The experiments were conducted in a RBD with 7 treatments and 3 replications.

Corresponding Author:**AY Munj**

Jr. Entomologist, Regional Fruit
Research Station, Vengurle
Maharashtra, India

Treatment details

T ₁	:	Foliar application of <i>Verticillium lecanii</i> @ 5 g/l (1x10 ⁸ cfu/g) – commercial product
T ₂	:	Foliar application of <i>Metarhizium anisopliae</i> (IIHR liquid formulation @ 1ml/10 l)
T ₃	:	Foliar application of consortia of <i>M. anisopliae</i> + <i>B. bassiana</i> (IIHR liquid formulation @ 1ml/10 l)
T ₄	:	Spray of IIHR formulation of <i>Metarhizium anisopliae</i> (IIHR oil formulation @ 0.5 ml/l)
T ₅	:	Spray of IIHR formulation of <i>Beauveria bassiana</i> (IIHR liquid formulation @ 1ml/l)
T ₆	:	1 st spray of spinosad 45 SL @ 0.004% at panicle emergence stage followed by 2 nd spray (21 days after 1 st) with thiamethoxam @ 0.008% and 3 rd need based spray of neemazal 10000 ppm @ 3ml/l (Standard Check)
T ₇	:	Control

Spray schedule

1 st spray -	At panicle initiation stage
2 nd spray -	7 days after 1st spray
3 rd spray -	7 days after 2nd spray
4 th spray -	At pea nut stage
5 th spray -	At marble stage

Observations on hopper population

Ten panicles were labeled randomly on each tree and the number of hoppers (nymphs and adults) observed on these panicles were counted. The pre-treatment observations were recorded 24 hours before each spray and the post treatment observations were recorded 7 days after each spray.

Results and Discussion

The pooled data (2015-16 to 2017-18) recorded on the efficacy of different treatments against mango hopper at different centers are presented in table 1. The pre count observations recorded a day before insecticidal application were statistically non-significant at all the centers. This indicates that the mango hopper population was uniform throughout the experimental area. The data recorded 7 days after last spray revealed that for management of mango hopper, the treatment T₆ (Standard check) was the most effective at all the centers except in Lucknow.

Among the different entomopathogen treatments used, treatment T₄ (Spray of IIHR oil formulation of *Metarhizium anisopliae* @ 0.5 ml/l) was found to be the most effective at Bengaluru and Periyakulam and significantly superior to all other treatments. Treatment T₄ recorded the least hopper count at Bengaluru (1.04 hopper/panicle) and Periyakulam (10.90 hopper/panicle) at 7 days after last spray as against 11.57 and 22.79 hoppers/panicle in untreated control, respectively.

Apart from standard check, treatment T₁ (Foliar application of *Verticillium lecanii* @ 5 g/l – commercial product) was found

to be the most effective for management of mango hopper at Lucknow, Paria and Vengurle and was significantly superior to the rest of the treatments at Lucknow and Paria, whereas, at Vengurle, it was at par with T₄. Treatment T₁ recorded the least hopper count at Lucknow (0.93 hopper/panicle), Paria (4.52 hopper/panicle) and Vengurle (0.80 hopper/panicle) as against 8.67, 10.72 and 5.53 hoppers/panicle in untreated control, respectively.

The yield data recorded under different treatments at different centers is presented in table 2. The data revealed that the maximum yield was recorded in treatment T₆ at Bengaluru (78.20 kg/tree), Paria (52.89 kg/tree), Periyakulam (66.17 kg/tree) and Vengurle (33.18 kg/tree). Treatment T₄ recorded the maximum yield at Bengaluru (71.45 kg/tree), Lucknow (57.16 kg/tree), Periyakulam (62.69 kg/tree) and Vengurle (31.39 kg/tree) and was significantly superior to rest of the treatments at Bengaluru, Lucknow and Periyakulam. However, at Vengurle, T₄ was at par with T₁. At Paria, treatment T₁ recorded the maximum yield (46.10 kg/tree) which was significantly superior to all other treatments.

The B:C ratio obtained under different treatments at different centers is presented in Table 2. The data revealed that, treatment T₄ recorded the maximum B:C ratio at Bengaluru (4.15), Lucknow (8.73), Periyakulam (2.29) and Vengurle (1.42). Whereas, at Paria the maximum B:C ratio was recorded under treatment T₅ (1.84) followed by T₁ (1.54).

These results are in close agreement with Srivastava^[15] who reported the efficacy of *Verticillium lecanii* against mango hopper in Uttar Pradesh. Gurav^[16] reported *V. lecanii* @ 10⁹ cfu/ml. as the most effective treatment with 93.34% mortality of mango hopper at Dapoli, Maharashtra during 2012. Turkhade^[17], who studied the combination effect of different entomopathogens, reported that the combination of *V. lecanii* + *M. anisopliae* + *B. bassiana* was effective against mango hopper at Dapoli, Maharashtra during 2014.

Table 1: Efficacy of different treatments against mango hopper (pooled data of 2015-16, 2016-17 and 2017-18)

Treatment	Hopper population/panicle									
	Bengaluru		Lucknow		Paria		Periyakulam		Vengurle	
	Pre count	After last spray	Pre Count	After last Spray	Pre count	After last spray	Pre count	After last spray	Pre count	After last spray
T ₁	6.00 (2.45)*	2.10 (1.45)	6.20 (2.48)	0.93 (0.64)	8.76 (2.96)	4.52 (2.13)	15.85 (3.98)	15.72 (3.97)	8.43 (3.06)	0.80 (1.30)
T ₂	5.86 (2.41)	2.74 (1.65)	6.93 (2.59)	3.93 (1.84)	8.31 (2.88)	6.67 (2.58)	17.02 (4.13)	13.61 (3.75)	7.87 (2.96)	1.58 (1.65)
T ₃	7.22 (2.68)	2.93 (1.71)	7.90 (2.74)	4.07 (1.96)	7.67 (2.76)	6.72 (2.59)	15.75 (3.97)	14.97 (3.89)	8.97 (3.13)	1.42 (1.61)
T ₄	6.66 (2.58)	1.04 (1.01)	7.42 (2.57)	4.23 (1.85)	7.99 (2.83)	5.67 (2.38)	16.19 (4.02)	10.90 (3.31)	8.42 (3.05)	0.96 (1.35)
T ₅	5.93 (2.43)	4.06 (2.01)	8.03 (2.83)	5.90 (2.43)	8.40 (2.89)	5.75 (2.40)	17.97 (4.23)	15.13 (3.91)	8.38 (3.04)	1.94 (1.74)
T ₆	7.33 (2.70)	0.85 (0.92)	7.43 (2.64)	5.63 (2.04)	9.06 (3.01)	3.64 (1.51)	15.91 (3.99)	5.37 (2.29)	8.02 (2.99)	0.46 (1.14)
T ₇	6.66	11.57	7.83	8.67	8.46	10.72	15.85	22.79	8.93	5.53 (2.44)

	(2.58)	(3.40)	(2.67)	(2.88)	(2.91)	(3.27)	(3.98)	(4.69)	(3.13)	
CD (0.5%)	N.S.	0.24	NS	0.20	N.S.	0.08	NS	0.40	N.S.	0.30

*indicates $\sqrt{n+1}$ transformed values

Table 2: Yield and B: C ratio recorded in different treatments (pooled data of 2015-16, 2016-17 and 2017-18)

Sr. No.	Treatment	Yield (kg/tree)									
		Bengaluru		Lucknow		Paria		Periyakulam		Vengurle	
		Yield	B:C ratio	Yield	B:C ratio	Yield	B:C ratio	Yield	B:C ratio	Yield	B:C ratio
1	T ₁	61.95	1.27	37.27	4.47	46.10	1.54	48.87	1.75	30.97	1.38
2	T ₂	57.60	3.23	29.63	2.02	34.14	1.01	51.62	1.89	28.17	1.23
3	T ₃	59.30	3.46	27.55	1.29	36.67	1.36	50.71	2.11	27.81	1.20
4	T ₄	71.45	4.15	57.16	8.73	39.75	1.10	62.89	2.29	31.39	1.42
5	T ₅	50.15	2.24	36.50	4.35	42.76	1.80	49.42	1.79	28.93	1.22
6	T ₆	78.20	3.36	53.67	4.37	52.89	2.60	66.17	2.52	33.18	1.50
7	T ₇	33.40	-	20.77	--	19.63	-	43.62	-	24.00	-
	CD (0.5%)	8.43	-	2.70		3.70		1.23	-	2.96	

Conclusion

From the overall results, it can be concluded that for eco-friendly management of mango hopper, the oil based formulation of *M. anisopliae* supplied by IIHR, Bengaluru was the most effective at Bengaluru, Lucknow, Periyakulam and Vengurle, whereas, *V. lecanii* (commercial product) was found to be the most effective at Paria.

Acknowledgement

This work was conducted with financial support from the Indian Council of Agricultural Research (ICAR) and the collaborative organizations (SAUs/ICAR Institutes) under the ICAR-All India Coordinated Research Project (AICRP) on Fruits. The authors wish to acknowledge the support of the ICAR and the collaborative organizations in providing the requisite research facilities.

References

- Butani DK. Save your crop from insect pests. Indian Horticulture. 1962; 6(1):6-7.
- Kumar D, Roy CS, Yazdani SS, Hamid SF, Khan ZR. Effect of some insecticides against hopper complex on mango. Pesticides. 1985; 19(11):42-43.
- Rahman SK, Kuldeep MA. Mango hoppers: Bio ecology and management- A review. Agricultural Reviews. 2007; 28:49-55.
- Chavan SA, Dalvi MB, Munj AY, Patil PD, Salvi BR. Mango Plant Protection, Ed. 1, Dr. B. S. Konkan Krishi Vidyapeeth, Dapoli 2009, 1-2.
- Munj AY, Sawant BN, Malshe KV, Dheware RM, Narangalkar AL. Survey of important foliage pests of mango from South Konkan region of Maharashtra. Journal of Entomology and Zoology Studies. 2019; 7(1):684-686.
- Godase SK, Patil PD, Bhole SR. Evaluation of some synthetic pyrethroides against hopper complex of mango, *Mangifera indica* L. Pestology. 2001; 25(4):29-30.
- Godase SK, Bhole SR, Shivpuje PR, Patil BP. Assessment of yield loss in mango due to mango hopper (*I. niveosparsus*). Indian Journal of Agriculture Sciences. 2004; 74(7):370-372.
- Singh M, Gupta D, Gupta R. Incidence and control of mango leaf hoppers in the lower hills of Himachal Pradesh. Pest Management and Economic Zoology. 2006; 14(1, 2):49-54
- Shukla RP. Strategies of integrated insect pest management in mango- Global scenario. Presented in: Global Conference on Augmenting Production and Utilization of Mango: Biotic and Abiotic Stresses. June, 2011 at CISH, Lucknow, 2011, 113.
- Ghosh SK. Sustainable management of mango hopper (*A. atkinsoni* and *I. niveosparsus*) by using insect growth regulator, buprofezin 25 SC. Uttar Pradesh Journal of Zoology. 2013; 33(2):119-128.
- Ramesh Babu S, Virendra Singh. Bioefficacy of Tolfenpyrad 15 EC against hopper complex in mango. Pest Management in Horticulture Ecosystems. 2014; 20(1):22-25.
- Thiruvani T, Ganesh Kumar M, Kuttalam S. Field evaluation of a new indigenous thiamethoxam 25% WG formulation against mango hopper. Pestology. 2014; 38(4):76-78.
- Munj AY, Sawant BN, Raut RA, Dheware RM, Salvi BR. Evaluation of insecticides for management of mango hopper, *Idioscopus niveosparsus* Leth. Journal of Entomology and Zoology Studies. 2018; 6(6):950-952.
- Munj AY, Halgekar NY, Salvi BR. Effective management module to minimize the crop loss in Alphonso mango due to mango hopper. Environment and Ecology. 2015; 33(4B):1969-1971.
- Srivastava RP, Tandon PL. Natural occurrence of two entomogenous fungi pathogenic to mango leaf hopper. Indian Journal of Plant Pathology. 1986; 4:121-123.
- Gurav PK. Evaluation of entomopathogenic fungi and botanicals for the management of mango hopper and thrips. M. Sc. (Agri.) thesis submitted to Dr. B. S. Konkan Krishi Vidyapeeth, Dapoli, Dist. Ratnagiri, M.S, 2012.
- Turkhade PD. Ecofriendly management of mango hopper *Idioscopus niveosparsus*. M. Sc. (Agri.) thesis submitted to Dr. B. S. Konkan Krishi Vidyapeeth, Dapoli, Dist. Ratnagiri, M.S, 2015.